

19  **Europäisches Patentamt**
European Patent Office
Office européen des brevets

11 Publication number:

0 119 448
A1

12

EUROPEAN PATENT APPLICATION

21 Application number: 84101392.3

51 Int. Cl.³: **C 07 H 21/00, C 07 H 19/00,**
G 01 N 33/48

22 Date of filing: 10.02.84

30 Priority: 14.02.83 JP 22516/83
 28.04.83 JP 75878/83

71 Applicant: **WAKUNAGA SEIYAKU KABUSHIKI KAISHA,**
 1-39 Fukushima 3-chome Fukushima-ku, Osaka-shi
 Osaka-fu (JP)

43 Date of publication of application: 26.09.84
 Bulletin 84/39


72 Inventor: **Miyoshi, Kenichi, 2521-17, Toge Kabe-Cho**
Asakita-Ku, Hiroshima-Shi Hiroshima-Ken (JP)
 Inventor: **Suzuki, Masanori, 208, Ku Koyo-Cho**
Asakita-ku, Hiroshima-Shi Hiroshima-Ken (JP)
 Inventor: **Fuwa, Toru, 6-17-802, Komachi Naka-Ku,**
Hiroshima-Shi Hiroshima-Ken (JP)

84 Designated Contracting States: **DE FR GB**


74 Representative: **Kraus, Walter, Dr. et al, Patentanwälte**
Kraus & Weisert Imgardstrasse 15,
D-8000 München 71 (DE)

54 **Oligonucleotide derivatives and production thereof.**

57 Biotin- or 2,4-dinitrophenyl-oligodeoxyribo-nucleotide derivative represented by the formula shown below and its preparation method is disclosed:



wherein:
 R is



m and n are each 0 or a natural number; R' is a straight or branched hydrocarbon residue; and B is a base constituting the nucleotide (in which when a plural number of B exist, they may be either identical or different). This compound is useful for use as a non-radioactive affinity probe for nucleic acid or a primer.

EP 0 119 448 A1

OLIGONUCLEOTIDE DERIVATIVES AND PRODUCTION THEREOF

BACKGROUND OF THE INVENTIONTechnical Field

5 This invention relates generally to oligonucleo-
tide derivatives. More specifically, the present in-
vention relates to an oligonucleotide derivative
comprising biotin or 2,4-dinitrobenzene bonded to a
moiety other than the base of the nucleotide. The
10 present invention also relates to a method for produc-
ing such an oligonucleotide.

Prior Art

 Biotin, which is a member of the vitamin B complex
and is also called vitamin H, is necessary for the
15 growth of many animals and vegetables. On the other hand,
biotin is known to interact potently with avidin in
egg white, and this characteristic is put into prac-
tical application by utilizing biotin in the form of
its derivative such as a biotin-avidin reagent. This
20 is applied as a biochemical reagent for measurement of
antigenic density per cell, radioimmunoassay, enzyme
immunoassay, etc. Also, a DNA probe having biotin
bonded to a nucleic acid has been developed and is now
commercially available for diagnosis of infectious and
25 hereditary diseases (Proc. Natl. Acad. Sci. USA, 78,
6633 - 6637, 1981). The biotin nucleotide derivative
in this DNA probe is prepared enzymatically with DNA
or RNA as the template by using the biotin derivative
of cytidine triphosphate in place of cytidine tri-
30 phosphate (dCTP).

 However, according to our knowledge, a biotin
nucleotide derivative thus prepared involves the
following problems.

 a) Since biotin is included at the base moiety
35 of the nucleotide, a change occurs in the melting
temperature (T_m value) inherent in the oligonucleotide
employed.

b) Synthesis of a cytosine derivative is difficult (in view of the above cited reference).

c) Synthesis of a DNA having a desired and predetermined base sequence is difficult.

5 For these reasons, biotin-nucleotide derivatives at the present stage have only a narrow scope of applications and are limited in their usefulness.

10 Meanwhile, Vincent et al have recently developed a DNA probe having 2,4-dinitrophenyl (hereinafter abbreviated as DNP) group bonded to nucleic acid (Nucl. Acids Res., 10, 6787 - 6796, 1982). They cause a DNP derivative of adenosine triphosphate (ATP) to be incorporated into a DNA strand to be hybridized with a DNA having a complementary base sequence and
15 then detect the objective DNA by adding successively a rabbit antiserum against DNP and peroxide-labelled goat antirabbit I_g. The DNA strand employed here is a fragment taken from nature.

20 However, according to our knowledge, the DNP-nucleotide derivative thus prepared involves the following problems.

a) Since DNP is included at the base moiety of the nucleotide, a change occurs in the melting temperature (T_m value) inherent in the oligonucleotide
25 employed.

b) Synthesis of a DNA having a desired and predetermined base sequence is difficult.

30 For these reasons, DNP-nucleotide derivatives at the present stage also have only a narrow scope of applications and are limited in their usefulness.



SUMMARY OF THE INVENTION

35 An object of the present invention is to overcome the drawbacks described above. This invention accomplishes this object by providing an oligonucleotide comprising biotin or 2,4-dinitrobenzene bonded to a specific oligodeoxyribonucleotide at a specific site


other than the nucleotide base.

The oligonucleotide derivative according to the present invention comprises an oligodeoxyribonucleotide represented by the formula [VII] set forth
5 below.

Further, the method for preparing the oligonucleotide derivative according to the present invention comprises causing biotin or 2,4-dinitrobenzene to be bonded to the terminal amino group of an oligo-
10 nucleotide derivative represented by the formula [VI] set forth below to obtain an oligodeoxyribonucleotide represented by the formula [VII] set forth below.



In the above formulae, R is



m and n are each 0 or any desired natural number; R^1 is a straight or branched hydrocarbon residue; and B is a base constituting the nucleotide. (When a plural
35 number of B exist, they may be either identical or different).

Advantageous Features

The biotin- or DNP-oligodeoxyribonucleotides

synthesized according to the present invention can circumvent the drawbacks of the non-radioactive affinity probe for nucleic acid as described above and has the following advantages.

- 5 a) Since neither biotin nor DNP is included at the base moiety of the nucleotide, it is stable without change in the melting temperature (T_m value).
- b) Synthesis of a biotin- or DNP-oligonucleotide having any base sequence is possible.
- 10 c) A short chain oligomer is sufficient for a probe.
- d) Synthesis is very simple, thus permitting large scale synthesis thereof, and storage for a long term is also possible.
- 15 e) It can also be utilized as a primer (DNA fragment in synthesis of template).

 Because of such advantages, the oligonucleotide derivatives may be considered to be utilizable for a wider scope of applications. More specifically, for
20 example, the biotin- or DNP-oligonucleotide is utilizable as a non-radioactive affinity probe for nucleic acid or as a primer as described above, and its detection may be performed according to various methods such as sedimentation by an antibody, activity
25 assay of an enzyme, visualization with an affinity column fluorochrome, etc. Furthermore, the oligonucleotide derivative of the present invention is advantageous with respect to the risk of exposure, cost, disposal of the waste and storability, as compared with
30 the radioactive probe (^{32}P).

 In addition, the DNP group can easily be detected with a commercially available rabbit antiserum (e.g., Code No. 61-006-1, produced by Miles Laboratories) or a monoclonal antibody against DNP.

35 BRIEF DESCRIPTION OF THE DRAWINGS

 FIG. 1 is a flow chart indicating an example of the method for synthesis of the compound of the present

invention in Examples 1 and 2 set forth hereinafter;

FIG. 2 is a flow chart of the method for synthesis of the compound of the present invention as illustrated in the Experiments in Examples 1 and 2;

5 FIGS. 3A through 3D are chromatograms indicating the results of the high speed liquid chromatography of the compounds [VI] as illustrated in the Experiments in Examples 1 and 2;

10 FIGS. 4 and 5 are charts showing elution patterns of the high speed liquid chromatography in Example 1;

FIGS. 6 and 7 are charts each showing the results of electrophoresis in Example 1; and


FIG. 8 is a chart showing elution patterns of the high-speed liquid chromatography in Example 2.

15 DETAILED DESCRIPTION OF THE INVENTION

Oligonucleotide derivative [VII]

The oligonucleotide derivative according to the present invention is represented by the formula [VII] as described above.

20 In the above formula, the symbol B is conventionally used for showing the deoxyribonucleoside residue after removal of 3'- and 5'-hydroxyl groups from 2'-deoxyribonucleoside, and has specifically the structure shown below:



30

The substituent B represents a base constituting the nucleotide and is usually adenine, thymine, cytosine or guanine. When a plural number of B exist in the compound [VII], they may be the same or different.

Each of m and n is 0 or a natural number. The polymerization degree of the oligonucleotide derivative

of the present invention is represented in terms of $\underline{m} + \underline{n}$ because the fractions with polymerization degrees of \underline{m} and \underline{n} , respectively, are condensed according to a preferred preparation method in the present invention (as hereinafter described in detail). In this case, \underline{m} may practically be 0 to 6, particularly 1 to 4, and \underline{n} may practically be 0 to 40, particularly 0 to 20.

The group R^1 is a divalent straight or branched hydrocarbon residue for coupling the nucleic acid moiety of the compound [VII] with the biotin or DNP moiety. This may suitably be a straight or branched alkylene group, particularly having about 2 to 20 carbon atoms. Preferably, R^1 is an alkylene group having 2 to 6 carbon atoms, a straight C_2-C_6 alkylene being more preferable.

Synthesis of the compound [VII]


The compound [VII], namely, the oligonucleotide derivative according to the present invention can be synthesized according to any desired method suited for the purpose.

A preferable method comprises causing biotin or 2,4-dinitrobenzene to be bonded to the amino group of an oligonucleotide derivative of the above formula [VI], namely, a compound in which a primary amino group has been introduced through the group R^1 into an oligodeoxynucleotide at its 5'-terminal phosphate group.


On the other hand, the compound of the formula [VI] can be synthesized according to the method which comprises synthesizing an oligonucleotide and introducing a primary amino group into the oligonucleotide formed so as to be extended from the 5'-hydroxyl group.

FIG. 1 shows a flow chart illustrating an example of the preferable synthetic method. The symbols in the flow chart have the meanings defined below (their significances or details being as described hereinafter).

- R^0 : a substituent for protection of a phosphoryl group, o-chlorophenyl group being generally used;
- R^1 : a divalent hydrocarbon residue;
- R^2 : a protective group for 5'-terminal hydroxyl group, dimethoxytrityl group being generally used;
- R^3 : a substituent which can easily be eliminated under the conditions under which all other protecting groups are stable to produce a phosphodiester, a cyanoethyl group being generally used;
- R^4 : a protecting group for amino group, trifluoroacetyl group being generally used;
- q : any desired natural number smaller than n ;
- m : 0 or any desired natural number;
- n : 0 or any desired natural number;
- B : a base;
- B' : a protected base, generally selected from N^6 -benzoyladenine, N -isobutyrylguanine, N^6 -benzoylcytosine and thymine (i.e., no protection needed);



 P : carrier through a spacer, usually one as shown below:

25




(wherein p is polymerization degree of styrene)

30



35



Synthesis of the compound [VI]

Generally speaking, the methods for synthesis of oligonucleotides are the triester method and the phosphite method, each involving the solid phase method and the liquid phase method. We have already established a technique for producing oligonucleotides according to the solid phase phosphotriester method. Our methods as disclosed in the references set forth below are preferred for synthesis of the compound [VI].

- Tetrahedron Letters 1979, 3635 (1979)
- Nucleic Acids Research 8, 5473 (1980)
- Nucleic Acids Research 8, 5491 (1980)
- Nucleic Acids Research 8, 5507 (1980)
- Nucleic Acids Research Symposium Series 7, 281 (1980)

As a method for introducing a primary amino group into the 5'-hydroxyl group of the oligonucleotide as synthesized above through a phosphate group, that is, as a method for synthesizing the compound [VI], there is our method as disclosed in Japanese Patent Application No. 138136/1982.

The method of synthesizing the compound [VI] according to an embodiment thereof is described below. As shown in Fig. 1, the derivative of the compound [I] from which the protecting group R^3 has been removed and the derivative of the compound [II] from which the protecting group R^2 has been removed are condensed, and these operations are repeated to synthesize the compound [III]. The method of synthesizing an oligonucleotide compound [III] is known in the art as mentioned above.

On the other hand, the compound of the formula [IV] is synthesized according to our method (reference: Japanese Patent Application No.138136/1982). More specifically, R^2 of the compound [I] is removed to form a 5'-hydroxyl compound, which is caused to react

with a phosphorylating agent (e.g., phosphoditri-
azolide, phosphodichloride or phosphodibenzotri-
azolide) to be phosphorylated and then condensed
with an amino alcohol compound of which amino group
5 is protected, namely, R^2-NH-R^1-OH [which compound
can be obtained by protecting the amino group of an
omega-aminoalcohol (NH_2-R^1-OH) with R^2], whereby
the compound [IV] can be obtained (reference: the
specification of the above patent application for
10 its details).

The compound [IV] after removal of the protect-
ing group R^3 therefrom is condensed with the compound
[III] from which the protecting group R^2 has been
removed to synthesize the compound [V]. Condensation
15 may be carried out according to a method which is
substantially the same as that for the condensation
in the synthesis of the compound [III].

By removing all of the protecting groups on the
compound [V] thus synthesized, the compound [VI] can
20 be obtained. The protecting group $CO-\text{---}P$, o-
chlorophenyl group and the acyl group in the base
moiety can be removed by treatment with a 0.5 M
tetramethylguanidine-pyridine-2-carboaldoxime in
dioxane-water (9 : 1, v/v) solution, followed by
25 alkali treatment (conc. ammonia water). In the
case where R^4 is trifluoroacetyl group, it can be
removed at the same time by treatment with
ammonia, but treatment with mercaptoethanol is
required in the case where R^4 is o-nitrophenylsul-
30 feryl group. When another group is employed for R^4 ,
still another treatment may be applicable under the
conditions where the oligonucleotide moiety is stable.
In this context, various methods have already been
known for synthesis of deoxyoligoribonucleotide, and
35 as to details concerning the kinds of protecting
groups, introduction or removal thereof and condensa-
tion other than those as mentioned above, reference

may be made to textbooks or review publications, for example, "Synthesis of Nucleosides and Nucleotides" (Maruzen, Japan, 1977), "Organic Chemistry of Nucleic Acids" (Kagaku Dojin, Japan, 1979), "Nucleic Acid" (Asakura Shoten, Japan, 1979), Tetrahedron, 34, 3143 (1978), Journal of the Society of Organic Synthetic Chemistry, Japan, 34, 723 (1978) and "Kagaku no Ryoiki" (Domain of Chemistry) 33, 566 (1979).

Synthesis of the compound [VII]

10 The oligodeoxyribonucleotide (the compound [VII]) can be prepared by bonding biotin or 2,4-dinitrobenzene to the primary amino group extended from the 5'-terminal of the above compound [VI].

(1) Reaction with biotin:

15 Bonding between the two compounds can be carried out according to any method which can realize formation of an amide bonding through dehydration between the carboxyl group of biotin and the amino group of the compound [VI]. When an amino group or a hydroxyl
20 group which can react with the carboxyl group of biotin exists in the compound [VI], the reaction can be carried out under appropriate protection of such groups. Thus, the expression "by bonding biotin to the terminal amino group of the oligonucleotide derivative represented by the formula [VI], the biotin-oligonucleotide
25 represented by the formula [VII] is obtained" is also inclusive of the case where the compound [VI] is protected as described above. Further, this expression is also inclusive of the case where biotin is in
30 the form of its functional derivative. Specific examples of functional derivatives of biotin are its acid halides and active esters.

 A preferable method for obtaining bonding between the amino group and biotin in this sense comprises a
35 reaction between the amino group of the oligonucleotide derivative and an active ester of biotin. An active ester of biotin is preferred because it can react

selectively only with the primary amino group extended from the 5'-terminal hydroxyl group without undergoing reaction with the amino group at the base moiety of the oligonucleotide and also because the reaction operation is simple. An "active ester of biotin" means a biotin derivative having an ester bonding which can readily react with other functional groups (usually amino groups), typically succinimide-, p-nitrophenyl-, benzotriazolide- and 2,4,5-trichlorophenyl-ester.

Another preferable method for causing bonding between the amino group and biotin comprises carrying out bonding of both in the presence of a condensing agent. Examples of suitable condensing agents are dicyclocarbodiimide, carbonylimidazole, and Woodward reagent "K".

According to any of these methods, the reaction method can be any one suited for the purpose. The specific reaction method for a given reaction system is suitably determined by referring to the experimental examples as given below and various textbooks such as "Peptide Synthesis" (Maruzen, Japan 1975) and "Chemistry of Proteins IV" (1977).

(2) Reaction with 2,4-dinitrobenzene:

The DNP-oligodeoxyribonucleotide (the compound [VII]) can be obtained by causing 2,4-dinitrobenzene to be bonded to the primary amino group extended from the 5'-terminal of the above compound [VI].

Bonding between these compounds can be accomplished according to any desired method which can realize formation of C - N bonding between the 1-position of 2,4-dinitrobenzene and the amino group of the compound [VI].

Bonding between the two compounds is generally accomplished through condensation with H - X elimination between the derivative of the former, namely DNP - X

(where X is a 1-substituent) and the amino group. X is preferably a halogen. The derivative wherein X is a halogen, namely, 1-halogeno-2,4-dinitrobenzene is preferred because it can react selectively only
5 with the primary amino group extended from the terminal 5'-hydroxyl group without undergoing the reaction with the amino group at the base moiety of the oligonucleotide. Above all, 1-fluoro-2,4-dinitrobenzene is readily available commercially,
10 and the reaction with the amino group of the compound [VI] can proceed under mild reaction conditions.

The reaction of a 1-halogeno-2,4-dinitrobenzene with the compound [VI] can be practiced in a homogeneous solution (the solvent being, for example, a
15 water-containing alcohol) or a heterogeneous solution (the solvent being, for example, water) of the two compounds at a temperature of the order of 10 to 50°C in the presence of an agent for capturing a hydrogen halide liberated (e.g., sodium hydrogen carbonate,
20 triethylamine, and potassium hydroxide). The desired product may be recovered by, for example, extraction. As to incorporation of DNP, reference may be made to an appropriate review publication, for example, "Course of Chemical Experiments 1, Chemistry of
25 Proteins II, p.118" (published by Maruzen Co., 1976).

EXPERIMENTAL EXAMPLES

Example 1 (preparation of biotin-oligonucleotide)

Flow chart:


The compound of the present invention (the compound (11) in FIG. 2) was prepared according to the flow
30 chart shown in FIG. 2.

In FIG. 2, the symbols have the following meanings.


B': N-Benzoyl-adenine

B : adenine

35 DMTr : dimethoxytrityl



5 5



(p: degree of
polymer-
ization)

R⁰: o-chlorophenyl

Et: ethyl

CE: cyanoethyl

10


m : 2

n' : 2

n : 12

R :

15



20

Synthesis of the compound [VI]:

Experiment 1-1

25 Dimethoxytrityladenosine/resin [(1)] (resin is merely a carrier, but since the desired compound carried on a resin is substantially the same in appearance as the resin itself, the compound carried on a resin is hereinafter called merely resin) in a
30 quantity of 300 mg (0.033 mmol), was washed three times with 10 ml of an isopropanol-methylene chloride (15:85, v/v) solution, and then the reaction (detritylation) was carried out four times each
35 for 5 minutes with 8 ml of a 1.0 M zinc bromide in an isopropanol-methylene chloride solution to obtain a resin [(2)]. The resin [(2)] was washed three times with 10 ml of an isopropanol-methylene chloride

solution, and after addition thereto of a solution of 150 mg of the dinucleotide [(3)] in pyridine, the mixture was subjected to azeotropic distillation to make the system anhydrous. Then, 150 mg (0.5 mmol) of mesitylenesulfonyl nitrotriazolide (hereinafter abbreviated MSNT) and 2 ml of anhydrous pyridine were added to the system, and the reaction (condensation) was carried out for 90 minutes. After the reaction, the reaction mixture was washed three times with 10 ml of pyridine, and 10 ml of a solution containing a catalytic amount (about 10 mg) of dimethylaminopyridine (hereinafter abbreviated DMAP) in acetic anhydride-pyridine (1:9, v/v) to carry out the reaction for 10 minutes, thereby protecting or masking the unreacted 5'-hydroxyl group through acetylation. The protected compound was washed with pyridine to obtain the compound [(4)'] (n=2). The above operation was repeated 6 times to obtain the compound [(4)] (n=12).

On the other hand, the reaction between 800 mg (0.71 mmol) 5'-hydroxy-dinucleotide [(5)] and o-chlorophenyl phosphoditriazolide was carried out in a solution of the latter in dioxane (1.0 mmol, 6 ml) for 2 hours, followed further by the reaction continued for 2 hours with addition of 300 mg (1.4 mmol) of trifluoroacetyl-6-aminohexanol and 115 mg (1.4 mmol) of 1-methyl-imidazole. After completion of the reaction, the solvent was evaporated, and the residue was dissolved in chloroform. Then, the solution was washed with water, a 0.5 M aqueous sodium dihydrogen phosphate solution, a saturated aqueous sodium hydrogen carbonate solution and an aqueous 5% sodium chloride solution, respectively, and dried over anhydrous sodium sulfate. The chloroform layer was concentrated and purified on a silica gel column, eluting with chloroform containing 0 to 4% methanol. The eluate was concentrated and added dropwise into pentane to obtain a

powdery compound [(6)].

To the compound [(7)] obtained by detritylation of 115 mg (3.45 μ mol) the compound as prepared above [(4)] (n=12) according to the same procedure as described above was added the compound [(6)] (0.04 mmol) which had been decyanoethylated by treatment thereof with 3 ml of a triethylamine-pyridine-water (1 : 3 : 1, v/v) solution. After the system was made anhydrous, 50 mg (0.2 mmol) of MSNT and 1 ml of pyridine were added thereto, and the reaction (condensation) was carried out for 90 minutes. After completion of the reaction, the product was washed with pyridine and methanol and dried to obtain a completely protected oligonucleotide derivatives [(9)].

15 mg of the oligonucleotide derivative [(9)] was added to 200 μ l of a solution of 0.5 M tetramethylguanidine-pyridine-2-carboaldoximate in dioxane-water (9 : 1, v/v), and reaction was carried out in a centrifugal precipitating tube at room temperature for 24 hours. After the reaction, conc. ammonia water (2.5 ml) was added thereto, and, with the vessel in sealed state, the reaction was carried out at 50°C overnight. After completion of the reaction, the mixture was filtered, and the filtrate was concentrated. The concentrate was dissolved in water and washed with ether. The aqueous layer was concentrated and purified by desalting on Sephadex G-50 (ϕ 1.5 x 120 cm, eluting with 0.05 M triethylammonium bicarbonate buffer, pH 7.5) to obtain a pentadecadenylic acid derivative [(10)].

According to similar procedures, the oligonucleotide derivatives of Experiments 1-2, 1-3 and 1-4 were prepared. The compounds as synthesized above are shown in Table 1.

Table 1

Deriva- tive	Identities of Compound (10)	
	m+n	(B) _{m+n} B
5 Experiment		
1-1	14	A A A A A A A A A A A A A A
1-2	14	T T T T T T T T T T T T T T
1-3	14	G G A T G C A T C A C C A C C
10 1-4	16	A A T C T G G T G A G A A G C G C

In the above Table, A represents adenine, T thymine, G guanine and C cytosine.

The results of high-speed liquid chromatography for these four types of compounds are shown in FIG. 3, in which A through D are charts for the compounds of Experiments 1-1 through 1-4, respectively.

Preparation of biotin-pentadecaadenylic acid [(11)]:

Experiment 2-1

20 About 1.0 OD of the pentadecaadenylic acid derivative [(10)] as prepared in the above experiment 1-1 was dissolved in 10 μ l of 0.1 M an aqueous sodium hydrogen carbonate solution (pH 8.3) and, with addition of 10 μ l of a solution of biotin succinimide ester in 25 dimethylformamide (corresponding to some hundred-fold excess), the reaction was carried out at 4°C overnight or at room temperature for 4 hours to synthesize biotin-pentadecaadenylic acid [(11)].

30 The reaction was monitored by high-speed liquid chromatography and 20% polyacrylamide gel electrophoresis.

The same operation as in Experiment 2-1 was also applied for the compounds [(10)] prepared in the above Experiments 1-2, 1-3 and 1-4, respectively, to prepare 35 the respective compounds [(11)]. The experiments conducted here are called Experiments 2-2, 2-3 and 2-4, respectively. The compounds prepared in Experiments 2 are shown in Table 2.

Table 2

Deriva- tive Experiment	Identities of Compound (11)	
	m+n	(B) _{m+n} ^B
2-1	14	A A A A A A A A A A A A A A
2-2	14	T T T T T T T T T T T T T T
2-3	14	G G A T G C A T C A C C A C C
2-4	16	A A T C T G G T G A G A A G C G C

In the above Table, A represents adenine, T thymine, G guanine and C cytosine.

Reference Experiment 3

For the purpose of comparison with respect to reactivity, the compounds [(12)] having 5'-hydroxyl group obtained through deprotection of the oligonucleotides [(7)] as synthesized above Experiments 1 were also similarly caused to react with biotin succinimide ester.

The reference experiments are called Experiments 3-1, 3-2, 3-3, 3-4, respectively. The compounds used in Experiments 3 are shown in Table 3.

Table 3

Deriva- tive Experiment	Identities of Compound (12)	
	n	(B) _n ^B
3-1	12	A A A A A A A A A A A A
3-2	12	T T T T T T T T T T T T
3-3	12	A T G C A T C A C C A C C
3-4	14	T C T G G T G A G A A G C G C

In the above Table, A represents adenine, T thymine, G guanine and C cytosine.

The above results are shown in FIG. 4 and FIG. 5 (results of high-speed liquid chromatography) and in FIG. 6 and FIG. 7 (results of electrophoresis).

FIG. 4 shows elution patterns of high-speed liquid chromatography. In this Figure, each of the numerals 1 shows the chromatogram of the compound before the reaction, each of the numerals 2 that of a mixture of the compounds before and after the reaction, and each of the numerals 3 that of the compound after the reaction with the biotin active ester. (a) is the chromatogram in Experiment 3-1, (b) is the chromatogram in Experiment 2-1, (c) is the chromatogram in Experiment 3-2, and (d) is the chromatogram in Experiment 2-2, respectively. The numerical value on the peak shows the retention time.

FIG. 5 shows elution patterns of high-speed liquid chromatography. In this Figure, each of the numerals 1 shows the chromatogram of the compound before the reaction, each of the numerals 2 that of a mixture of the compounds before and after the reaction, and each of the numerals 3 that of the compound after the reaction with biotin active ester. (e) is the chromatogram in Experiment 3-3, (f) is in Experiment 2-3, (g) is in Experiment 3-4, and (d) is in Experiment 2-4, respectively. The numerical value on the peak shows the retention time.

FIG. 6 shows the results of electrophoresis, in which (a) and (b) show the electrophoresis profile before and after reaction in Experiment 3-1, respectively. Similarly, (c) and (d) show Experiment 2-1, (e) and (f) show Experiment 3-2, (g) and (h) show Experiment 2-2, respectively, with the biotin active ester. XC indicates the band of xylene cyanol and BPB that of bromophenol blue, and both are used as the labels in electrophoresis. In this drawing, the upper side indicates the minus side, and the lower side the plus side.

FIG. 7 shows the results of electrophoresis, in which (i) and (j) show Experiment 3-3, (k) and (l) show Experiment 2-3, (m) and (n) show Experiment 3-4, (p) and (o) show Experiment 2-4, respectively, with the biotin active ester. BPB has the same meaning as mentioned above, and in this drawing, the upper side indicates the minus side, and the lower side the plus side.

As seen from the results of high-speed liquid chromatography (FIG. 4 and FIG. 5), in Experiment 3, the compound having a 5'-hydroxyl group represented by the formula (12) (FIG. 4a-1,3, FIG. 4c-1,3, FIG. 5e-1,3 and FIG. 5g-1,3 did not react with biotin, and gave a peak at the same position before and after reaction. In contrast, in Experiment 2, the oligonucleotide derivative [formula (10)] reacted with biotin and yielded a new peak. The peak of the starting material (FIG. 4b-1, FIG. 4d-1, FIG. 5f-1 and FIG. 5h-1) disappeared, and a new peak through the reaction with the biotin active ester was revealed (FIG. 4b-3, FIG. 4d-3, FIG. 5f-3 and FIG. 5h-3). In FIG. 4 to FIG. 5, each of the numerals 2 shows the chromatogram of a mixture of the compounds before and after the reaction. These results clearly suggest that the compounds [(12)] never react with the biotin active ester, and that the compounds [(10)] react quantitatively to give the compounds [(11)].

Then, electrophoresis analysis of the reaction between the 5'-hydroxyl compound [(12)] and the biotin active ester in FIG. 6 and FIG. 7 [see (a)/(b), (e)/(f), (i)/(j) and (m)/(n)] shows that no difference can be seen in the positions of the bands before the reaction [(a), (e), (i) and (m)] and those after the reaction [(b), (f), (j) and (n)]. It can thus be concluded that no reaction with biotin occurred. On the other hand, in the reaction of the oligonucleotide [(10)] having primary amino group with the biotin active

ester [(c)/(d), (g)/(h), (k)/(l) and (o)/(p)] in FIGS. 6 and 7, the positions of the bands before the reaction [(c), (g), (k) and (o)] are different from those after the reaction [(d), (h), (l) and (n)], thus
 5 indicating that the reaction with biotin occurred.


From the above results, it can be seen that the compound having primary amino group as prepared above reacts selectively and quantitatively with the biotin active ester.

10 Example 2 (preparation of DNP-oligonucleotide)

Flow chart:

The compound of the present invention (the compound [(11)] in the Figure) was produced according to the flow chart in FIG. 2.

15 The symbols in FIG. 2 are the same as those described in Example 1, except for the following:



20

Synthesis of the compound [VI] ([(10)] in FIG. 2):
Experiment 1-1

The compound [VI] was prepared in the same manner as described in Example 1.

25 Preparation of 2,4-dinitrophenyl-pentadecaadenylic acid [(11)]:
Experiment 2-1

About 1.0 OD of the pentadecaadenylic acid derivative [(10)] synthesized in the above Experiment 1-1
 30 was dissolved in 0.1 M sodium bicarbonate solution (pH 8.3), and the reaction was carried out with addition of 5 μ l (great excess) of a solution of 1-fluoro-2,4-dinitrobenzene in ethanol (50 mg/ml) at 37°C for 2 hours. Then, the product was diluted with 30 μ l of water
 35 and washed four times with 150 μ l of ether to obtain 2,4-dinitrophenyl-pentadecaadenylic acid [(11)]. The reaction was monitored by high-speed liquid chromatography.

The compounds [(10)] synthesized in the above Experiments 1-2, 1-3 and 1-4 were also subjected to the same operation as described in Experiment 2-1 to prepare compounds [(11)] corresponding to respective compounds. The experiments conducted here are called Experiments 2-2, 2-3 and 2-4.

The compounds prepared in Experiment 2 are shown in Table 4. This Table is the same as Table 2 in Example 1.

10

Table 4

Deriva- tive Experiment	Identities of Compound (11)	
	m+n	(B) _{m+n} ^B
2-1	14	A A A A A A A A A A A A A A A A
2-2	14	T T T T T T T T T T T T T T T T
2-3	14	G G A T G C A T C A C C A C C C
2-4	16	A A T C T G G T G A G A A G C G C

20

In the above Table, A represents adenine, T thymine, G guanine and C cytosine.

Reference Experiment 3

For the purpose of comparison with respect to reactivity, the compounds [(12)] having 5'-hydroxyl group obtained by deprotection of the oligonucleotides [(7)] as prepared above Experiments 1 were also allowed to react in the same manner with 1-fluoro-2,4-dinitrobenzene.

The reference experiments conducted here are called Experiments 3-1, 3-2, 3-3, 3-4. The compounds used in Experiments 3 are shown in Table 5. This Table is the same as Table 3 in Example 1.

35

Table 5

Deriva- tive	Identities of Compound (12)	
	n	(B) _n ^B
Experiment		
3-1	12	A A A A A A A A A A A A
3-2	12	T T T T T T T T T T T T
3-3	12	A T G C A T C A C C A C C
3-4	14	T C T G G T G A G A A G C G C

In the above Table, A represents adenine, T thymine, G guanine and C cytosine.

The above results are shown in FIG. 8 (results of high-speed liquid chromatography).

FIG. 8 shows elution patterns of high-speed liquid chromatography. In this Figure, each of the numerals 1 indicates the chromatogram of the compound before the reaction, and each of the numerals 2 indicates that of the compound after the reaction with 1-fluoro-2,4-dinitrobenzene. (a) shows the chromatogram in Experiment 3-1, (b) in Experiment 2-1, (c) in Experiment 3-2, (d) in Experiment 2-2, (e) in Experiment 3-3, (f) in Experiment 2-3, (g) in Experiment 3-4, and (h) in Experiment 2-4, respectively. The numerical value on the peak shows the retention time.

As can be seen from these results, the compounds having 5'-hydroxyl group represented by the formula [(12)] (a-1, c-1, e-1 and g-1 in FIG. 8) do not react with 1-fluoro-2,4-dinitrobenzene (a-2, c-2, e-2 and g-2 in FIG. 8).

In contrast, the oligonucleotide derivative [(10)] reacted with 1-fluoro-2,4-dinitrobenzene, and yielded a new peak. The peak of the starting material (b-1, d-1, f-1 and h-1 in FIG. 8) disappeared and a new compound is formed through the reaction (b-2, d-2, f-2 and h-2).

In other words, it can be seen that the compound

[(10)] having a primary amino group reacts selectively and quantitatively with 1-fluoro-2,4-dinitrobenzene to give the compound [(11)] and that the compound [(12)] never reacts with it.

5 Each of the peaks obtained upon elution at a retention time of about 5 minutes in FIG. 8 may be considered to correspond to 2,4-dinitrophenol.

10 In Examples 1 and 2, high-speed liquid chromatography was conducted by means of HPLC System Tri-Roter III produced by Nippon Bunko under the following measurement conditions:

Column: μ - Bondapak C18 (Waters)


Flow rate: 2 ml/min.

15 Eluant: 20 mM-TEAA buffer (pH 7.2) containing acetonitrile


Concentration gradient: conc. of acetonitrile 6 to 14%/16 min. (15% after 16 min.)

WHAT IS CLAIMED IS:

1. A nucleotide derivative, comprising an oligo-nucleotide represented by the following formula [VII]:



wherein: R is




m and n are each 0 or a natural number; R^1 is a straight or branched hydrocarbon residue; and B is a base constituting the nucleotide (in which when a plural number of B exist, they may be either identical or different).

2. A nucleotide derivative according to Claim 1, wherein the base B is selected from the group consisting of adenine, thymine, cytosine and guanine.


3. A nucleotide derivative according to Claim 1, wherein R^1 is a straight or branched alkylene group having 2 to 20 carbon atoms.

4. A nucleotide derivative according to Claim 1, wherein m is a natural number of 0 to 6, and n is a natural number of 0 to 40.



5. A nucleotide derivative according to Claim 1, wherein R is



6. A nucleotide derivative according to Claim 1, wherein R is





7. A method for preparing a biotin nucleotide derivative, which comprises causing biotin to be bonded to the terminal amino group of an oligonucleotide derivative represented by the formula [VI] shown below to obtain a biotin-deoxyribonucleotide represented by the formula [VII] shown below:



wherein: \underline{m} and \underline{n} are each 0 or a natural number; R^1 is a straight or branched hydrocarbon residue; and B is a base constituting the nucleotide (in which when a plural number of B exist, they may be either identical or different).

8. A method according to Claim 7, wherein the bonding between the amino group and biotin is accomplished through the reaction of the amino group with an active ester of biotin.
9. A method according to Claim 8, wherein the active ester of biotin is biotinsuccinimide or biotin-p-nitrophenyl ester.
10. A method according to Claim 7, wherein the bonding between the amino group and biotin is accomplished under the action of a condensing agent.
11. A method according to Claim 10, wherein the condensing agent is dicyclohexylcarbodiimide.
12. A method for preparing a 2,4-dinitrophenyl-nucleotide derivative, which comprises causing 1-halo-2,4-dinitrobenzene to be bonded to the terminal amino group of an oligonucleotide derivative represented by the formula [VI] shown below through dehydrohalogenation to obtain a 2,4-dinitrophenyl-deoxyribo-nucleotide represented by the formula [VII] shown below:



wherein: \underline{m} and \underline{n} are each 0 or any desired natural number; R^1 is a straight or branched hydrocarbon residue; and B is a base constituting the nucleotide (in which when a plural number of B exist, they may be either identical or different).

0119448

27

13. A method according to Claim 12, wherein the 1-halogeno-2,4-dinitrobenzene is 1-fluoro-2,4-dinitrobenzene.

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6





FIG. 7






FIG. 8



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
Y	CHEMICAL & PHARMACEUTICAL BULLETIN, vol. 28, no. 6, 1980, pages 1695-1703, Pharmaceutical Society of Japan SHIGEKO NISHIMURA et al.: "Synthetic nucleosides and nucleotides. XV. 5-Dimethylamino-2-oxidoisoquinolin-1-yl diazomethane: a novel water-soluble fluorescent labeling agent for nucleotides" * Page 1695; page 1698, ref. 18 *	1	C 07 H 21/00 C 07 H 19/00 G 01 N 33/48
Y	--- CHEMICAL ABSTRACTS, vol. 94, no. 23, 8th June 1981, page 241, no. 187647x, Columbus, Ohio, US W.J. SMAGOWICZ et al.: "New substrate analogs for RNA polymerase" & STUD. BIOPHYS. 1980, 81(2-3), 135-6 * Abstract *	1	
Y	--- BIOCHEMISTRY, vol. 20, no. 19, 15th September 1981, pages 5538-5546, American Chemical Society, W.J. SMAGOWICZ et al.: "Properties of P3 esters of nucleoside triphosphates as substrates for RNA polymerase from Escherichia coli" * Pages 5538-5539 *	1	C 07 H 21/00 C 07 H 19/00
	--- -/-		
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 22-05-1984	Examiner VERHULST W.
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			



DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y	NUCLEIC ACIDS SYNTHESIS: APPLICATIONS TO MOLECULAR BIOLOGY AND GENETIC ENGINEERING, series no. 7, 1980, pages 167-171, IRL Press Ltd., London, GB R.I. GUMPORT et al.: "T4 RNA ligase as a nucleic acid synthe- sis and modification reagent" * Pages 167-168 *	1	
Y	--- EP-A-0 063 879 (YALE UNIVERSITY) * Pages 1-13 *	1	
Y,D	--- CHEMICAL ABSTRACTS, vol. 96, no. 7, 15th February 1982, page 207, no. 47771z, Columbus, Ohio, US P. LANGER et al.: "Enzymic syn- thesis of biotin-labeled polynucleotides: novel nucleic acid affinity probes" & PROC. NATL. ACAD. SCI. U.S.A., 1981, 78(11), 6633-7 * Abstract *	1	TECHNICAL FIELDS SEARCHED (Int. Cl. 7)
Y,D	--- CHEMICAL ABSTRACTS, vol. 98, no. 9, 28th February 1983, abstract, no. 68363z, Columbus, Ohio, US C. VINCENT et al.: "Synthesis of 8-(2-4 dinitrophenyl 2-6 aminol-amino-adenosine 5' triphosphate: biological proper- ties potential uses" & NUCLEIC ACIDS RES. 1982, 10(21), 6787-96 * Abstract *	1	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 22-05-1984	Examiner VERHULST W.
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			